



The Anatomy of Developmental Predictors of Healthy Lives Study (TADPOHLS)

Margaret L. Kern, Lizbeth Benson, Emily Larson, Christopher B. Forrest, Katherine B. Bevans & Laurence Steinberg

To cite this article: Margaret L. Kern, Lizbeth Benson, Emily Larson, Christopher B. Forrest, Katherine B. Bevans & Laurence Steinberg (2015): The Anatomy of Developmental Predictors of Healthy Lives Study (TADPOHLS), *Applied Developmental Science*, DOI: [10.1080/10888691.2015.1095642](https://doi.org/10.1080/10888691.2015.1095642)

To link to this article: <http://dx.doi.org/10.1080/10888691.2015.1095642>

 [View supplementary material](#) 

 Published online: 04 Nov 2015.

 [Submit your article to this journal](#) 

 Article views: 10

 [View related articles](#) 

 [View Crossmark data](#) 

The Anatomy of Developmental Predictors of Healthy Lives Study (TADPOHLS)

Margaret L. Kern

*University of Pennsylvania
The University of Melbourne*

Lizbeth Benson

*University of Pennsylvania
Pennsylvania State University*

Emily Larson

University of Pennsylvania

Christopher B. Forrest and Katherine B. Bevans

The Children's Hospital of Pennsylvania

Laurence Steinberg

*Temple University
King Abdulaziz University*

Numerous studies have followed people across significant portions of their lives. Secondary analyses with these studies offer opportunities to study life trajectories across diverse samples. To aid integrative efforts, we introduce The Anatomy of Developmental Predictors of Healthy Lives Study (TADPOHLS), a data base that categorizes items and constructs from 14 prospective longitudinal studies that followed participants from adolescence into adulthood. To classify items and measures, we created an extensive typology that provides a common language for categorizing study concepts. We illustrate the utility of the data base by examining adolescent perseverance and optimism as predictors of physical health outcomes across six studies. Adolescent perseverance and optimism were related to better physical health outcomes 15 to 20 years later. Overall, the data base offers a resource that contributes toward life-span studies of positive psychological and physical health.

“Long term longitudinal studies are like mature trees . . . like a century-old oak, such studies are rare resources and can add to our knowledge base in ways newer longitudinal studies cannot.”

Daniel Mroczek (2014)

Longitudinal studies are invaluable for investigating the development of health and well-being over the life course. There are now numerous studies available for secondary data analyses that have followed participants across decades of their lives. These investigations have collected detailed information on personal factors, environments, behaviors, physical health, and psychological functioning. Analysts wishing to use one of these datasets can feel overwhelmed in finding and selecting the right one for their purpose. To address this challenge, we developed The Anatomy of Developmental

Address correspondence to Margaret L. Kern, Melbourne Graduate School of Education, The University of Melbourne, Melbourne, VIC 3010, Australia. E-mail: Margaret.Kern@unimelb.edu.au

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/hads.

Predictors Of Healthy Lives Study (TADPOHLS) data base, which provides a typology of items and constructs from 14 longitudinal studies.

Longitudinal studies are rich sources of data that potentially can be used to examine temporal associations and change, control for confounding covariates, and capture different contexts of development (Hofer, Berg, & Era, 2003; Hofer & Sliwinski, 2001; Salthouse & Nesselroade, 2002; Slavich & Irwin, 2014). Each longitudinal study represents a major investment of time, money, and resources for researchers, participants, and funding agencies; therefore, it is incumbent on the research community to make optimal use of these data. However, any single longitudinal study is limited in numerous ways, including selection effects, historical period, attrition, missing data, and the quality of measures and documentation (Curran & Hussong, 2009; Hofer & Piccinin, 2009; Hofer & Sliwinski, 2006; Salthouse & Nesselroade, 2002).

Both cross-sectional and longitudinal designs have strong underlying assumptions that affect conclusions about development (Schaie, 1965). Cross-sequential, measurement burst, and other designs that combine elements of cross-sectional and longitudinal data have been developed to address some of the problems and limitations inherent to either type (e.g., Hofer & Sliwinski, 2001; Nesselroade, 2004; Salthouse & Nesselroade, 2002; Schaie, 1965; Schaie & Strother, 1968). Although we cannot change the structure of existing longitudinal studies, there is growing evidence that it is possible to directly integrate some studies together in a quasi-cross-sequential design to test developmental theories of health and well-being (e.g., Friedman, Kern, Hampson, & Duckworth, 2014; Hofer & Piccinin, 2010; Hussong, Curran, & Bauer, 2013; Kern, Hampson, Goldberg, & Friedman, 2014; Piccinin & Hofer, 2008). However, such work is neither straightforward nor simple.

Hofer and Piccinin (2009) summarized several levels of strategies that can be employed to build a comprehensive understanding of development with longitudinal studies. A typical approach is *sequential independent replication*, in which one study finds an association that is then replicated and extended in other studies. This approach is a key foundation of causal theory in contemporary social science. However, two common problems arise. First, due to study limitations, differing methodologies across studies, and an overreliance on significance rather than effect sizes, a subsequent study that purports to address the same hypothesis as a predecessor may produce findings that are quantitatively or qualitatively different. For example, comparing multiple cross-sectional cohorts with the same individuals measured over time, Schaie and Strother (1968) found strikingly different patterns, suggesting that

developmental differences were more a function of cohort than developmental change. Second, due to the tendency within the field to focus on “new” findings, replications are often not conducted or have been only conceptual in nature, such that the self-correcting process of science has not occurred (Duncan, Engel, Claessens, & Dowsett, 2014; Ioannidis, 2012; Makel, Plucker, & Hegarty, 2012; Pashler & Harris, 2012).

Meta-analyses are often considered a gold standard for summarizing effects. In a typical meta-analysis, a comprehensive literature review is conducted based on a specified set of search criteria. Effects are standardized and combined to provide an average overall effect, and moderators of the effect can be examined. Several guidelines have been developed to regulate and evaluate the quality of reviews (e.g., Higgins & Green, 2011; Higgins et al., 2013; Moher, Liberati, Tetzlaff, & Altman; The PRISMA Group, 2009; Shea et al., 2007). Meta-analyses offer the opportunity to find commonalities across studies and help the field to become more unified (Staats, 1999). However, studies are often excluded from an analysis because effect sizes cannot be calculated from the statistics reported in the study. As non-significant findings are often not published, averaged effects can be overestimated. Furthermore, a common question is the extent to which it is even appropriate to combine the effect sizes. Divergent constructs may have the same label (a “jingle” fallacy) and similar constructs may have different names (a “jangle” fallacy) (Block, 1995; Peck, 2004). Thus, the analyses combine apples and oranges, creating more of a mixed fruit salad than a blended apple pie.

There is a growing body of literature focused on *mega-analysis* or *individual participant data analyses* (IDA). Rather than using the effect sizes reported in studies, IDA compiles the raw data from multiple studies, examines items and constructs for conceptual and structural overlap, combines data at the individual participant level, and then tests specific theoretical models using the larger pooled dataset. To establish comparability across studies, the investigator must go beyond construct labels and engage with the specific items and measures used in each study. Heterogeneity across studies can be directly included in the analysis, testing the boundaries of generalizability, rather than being treated as problematic noise (Curran & Hussong, 2009).

The IDA approaches have been used for years in medicine, genetics, and economics. Many of these studies harmonize variables across studies by finding a common metric (e.g., dichotomized items), and then directly combining the aligned data. As many clinical trials are registered before the study begins and the same measures are often used across studies, combining studies is a relatively straightforward process. Stewart

and Clarke (1995) provided practical methodology for such mega-analytic reviews, noting that the statistical component is often the least time consuming and easiest aspect of the project.

The IDA approaches have only recently appeared in the social sciences, where it is considerably more challenging to find commonalities across studies. Many of the existing longitudinal studies have included numerous constructs, items, and variables, but the questions depended upon the original investigators' interests. Even when similar constructs exist across studies, the wording of the items often differs in terms of temporal orientation (e.g., past seven days versus past 12 months), and the response options vary. Still, a growing number of studies suggest that IDA in the social sciences is indeed possible (Hussong et al., 2013). For example, in the Healthy Ageing Across the Life Course (HALCyon) research program, physical capabilities data were harmonized across eight UK studies to examine cross-sectional age and gender differences in the measures (Cooper et al., 2011). Bath, Deeg, and Poppelaars (2010) harmonized 26 variables from the Longitudinal Aging Study Amsterdam (LASA) in The Netherlands and the Nottingham Longitudinal Study of Activity and Ageing (LSAA) in the UK, including demographic composition, physical and mental health, physical activity, religious attendance, pet ownership, and health service utilization. These studies demonstrate that integrating data across studies at the item level is possible, provides greater power to test complex models, and allows direct comparisons of study heterogeneity.

Hofer and Piccinin (2009) described the benefits of coordinating research across multiple longitudinal studies to enable such integrative data analysis to occur. Collaborative efforts allow more detailed analyses to be done, improve detailed data checking across studies, promote appropriate analyses and more balanced interpretations of results, and allow wider endorsement and dissemination of results (Hussong et al., 2013; Stewart & Clarke, 1995). The Integrative Analysis of Longitudinal Studies on Aging (IALSA), a collaborative network of longitudinal studies on cognition, health, and personality, provides one of the best examples of such coordinated efforts, and has pioneered numerous strategies for aligning and combining studies (Hofer & Piccinin, 2009; Piccinin & Hofer, 2008). To date, about 100 studies have agreed to be a part of the network (see <https://www.maelstrom-research.org/mica/network/ialsa>).

To complement the IALSA and other such consortium resources, we introduce The Anatomy of Developmental Predictors Of Healthy Lives Study (TADPOHLS), a collection of studies that prospectively followed participants from adolescence into adulthood,

and included measures of psychological, social, and physical health at multiple measurement occasions. Our goal was to identify and classify overlapping studies that potentially can be integrated together to study healthy development from adolescence into adulthood. In creating our structure, we were inspired by the IALSA, and thus modeled our structure after the networks' early work. TADPOHLS adds a catalogue of studies that included assessments of physical and psychological health in both adolescence and adulthood, to enable the assessment of developmental trajectories of health and well-being as youth transition into adulthood. In addition, we contribute an extensive coding typology, which is particularly detailed in terms of physical health variables. The typology provides a common language for categorizing study concepts, allowing analysts to examine concepts both within and across studies.

In this article, we provide background information on the rationale and methodology for developing TADPOHLS, introduce the data base and typology, and describe the studies and information included. We then provide a simple illustrative example in which we used the data base to identify six overlapping studies, and combined data from these studies to examine prospective associations between perseverance and optimism in adolescence and physical health outcomes in adulthood.

DEVELOPING THE TADPOLHS DATA BASE

Study Identification

A key goal of the project was to aggregate studies with measures of both physical health and psychological well-being in adolescence and adulthood. To identify studies, we built upon Friedman et al. (2014), who identified 88 different longitudinal studies that included personality and health variables from different periods of the life span. The authors searched a set of existing data bases (e.g., the Henry Murray Research Archives, the Inter-University Consortium for Political and Social Research), as well as PsycInfo, and Google Scholar, using the keywords "personality," "health," and "longitudinal." We used the list of 88 studies as a foundation, selecting studies that included measurement occasions in both adolescence and adulthood. In addition, we searched the set of existing data bases (see Table 1 in Friedman et al., 2014) for additional studies that fit our inclusion criteria. Through this process, we identified 60 potential studies for further consideration.

Research assistants then attempted to track down the identified studies. To be included in the data base,

studies had to have (1) accessible, readable codebooks; (2) data that could be acquired by researchers; (3) measurement occasions that occurred in both adolescence (age 13–18) and adulthood (18+); and (4) measures that included items on physical health and positive psychological functioning.

Coding Typology

To classify item-level and construct-level concepts across studies, we created a comprehensive coding typology (see Supplemental Material 1). To frame our categories, we began with the well-known International Classification of Functioning (ICF), which is divided into three components: functioning, personal factors, and environmental factors. We divided functioning into physical and psychological components, and the latter we further divided into positive and negative components to reflect our interest in both psychological well-being and emotional distress. Lastly, we added behaviors to provide a more comprehensive set of health predictors. This step resulted in six broad categories: physical functioning, positive psychological functioning, negative psychological functioning,

health behaviors, individual differences, and socio-environmental factors.

As illustrated in Figure 1, we structured the typology hierarchically. Within each of the six categories, we defined specific outcomes (e.g., cardiovascular system, depression, externalizing behaviors, personality, physical activity, relationships, socioeconomic status), and sub-outcomes (e.g., blood pressure, depression diagnosis, bullying, Big Five Inventory, leisure time activities, teacher connectedness, education).

Coding Procedure

For each study, the TADPOHLS typology was used as a guiding manual for selecting and coding relevant items from the original study codebooks. Each selected item was coded into a Microsoft Access data base. For example, starting with the question “I wish I had more friends,” we first determined that the item was relevant to the positive psychological functioning category, and then recognized that it focused on relationships (an outcome). As the question dealt with lack of social connection, it was classified as “loneliness”

TABLE 1
Study Descriptions

<i>Abbr</i>	<i>Study Name</i>	<i>Start Year</i>	<i>Country</i>	<i>Years Follow Up</i>	<i>Measure Occasions</i>	<i>Baseline N</i>	<i>Age at Baseline</i>	<i>Sample Type</i>
AHCE	Adolescent Health Care Evaluation Study	1984	US	6	4	2,788	13–18	Specific characteristic
ESDS	British Birth Cohort Study: 1958 Cohort	1958	UK	55	10	16,000	Birth	Nationally representative
ESD2	British Birth Cohort Study: 1970 Cohort	1970	UK	34	6	17,415	Birth	Nationally representative
FTP	Family Transitions Project	1989	US	20	4	451	12	Specific population
HLSU	Harlem Longitudinal Study of Urban Black Youth	1968	US	26	5	668	12–18	Nationally representative
NLSA	National Longitudinal Study of Adolescent Health	1994	US	14	4	15,701	10–18	Nationally representative
NLS2	National Longitudinal Survey of Youth – 1997 Cohort	1997	US	14	12	8,984	12–17	Nationally representative
NLSC	National Longitudinal Survey of Youth – 1979 Child and Young Adult	1986	US	10	12	5,255	Birth - 22	nationally representative
NLSY	National Longitudinal Survey of Youth – 1979 Cohort	1979	US	27	23	12,686	14–22	Nationally representative
NSHD	British Birth Cohort Study: 1946 Cohort	1946	UK	59	21	5,362	Newborn	Convenience sample
TBSS	The Beginning School Study	1982	US	20	9	790	1st grade	Community representative
TLCS	Terman Life-Cycle Study of Children with High Ability	1921	US	90	15	1,528	3–19	Specific population
WCFA	Welfare, Children, and Families: A Three City Study	1999	US	7	3	2,402	0–4; 10–14	Specific population
YTP	Youth in Transition Project	1966	US	4	4	2,213	15–16	Nationally representative

Note. See Supplemental Material 2 for a more detailed overview of each study.




FIGURE 1 Typology hierarchical structure. Variables were coded into six major categories (top). Example outcomes, sub-outcomes, and items are given. See Supplemental Material 1 for full typology.

(a sub-outcome). This process was repeated for each question from each of the studies.

Over a two-year period, our team spent thousands of hours coding items into the data base, identifying over 8,000 items. To ensure inter-rater reliability between coders, the coders went over each other's work and met weekly to address any discrepancies. After all items were entered, four research assistants cleaned the data base, fixing any remaining inconsistencies, checking for spelling mistakes, wrong categories, and repetitious questions, and adding missing information.

Resulting Resource

Items were classified for 25 studies. We focus here on 14 studies in which we established data-sharing agreements and were able to obtain the data. Table 1 summarizes descriptive information about each study, and Supplemental Material 2 provides more detailed study descriptions. Studies were conducted in the United States and the United Kingdom. The earliest study (the Terman Life Cycle Study) began in 1921 and followed people across their entire lives; the latest study began in 2002. Baseline age ranged from prenatal to 18 years, sample sizes ranged from 451 to 17,415 individuals, and measurement occasions ranged from three to

23. The final data base includes 8,447 items. Table 2 summarizes the number of items available for each outcome. The full data base can be accessed through www.margaretkern.org/TADPOHLS.html.¹

DISCUSSION

The TADPOHLS data base has classified items from longitudinal studies according to an extensive coding scheme. Although the goal of this data base is to enable integrative research and collaborative work, a wealth of research has already occurred using these data sets, as evidenced by the thousands of publications that other researchers have built their careers upon. We hope that the data base will help researchers identify studies that have similarities, making it easier to develop cross-study collaborations. The typology provides a structure for classifying items and measures across six broad areas. The detailed physical health categorization is a particularly useful addition.

¹To protect participants and honor data sharing agreements, the website provides detailed information about the variables available, and provides contact information or websites for each study, but does not provide the actual questionnaires, codebooks, or data. It is the user's responsibility to work directly with the original study investigators to obtain the data.

TABLE 2
Outcomes Included in Each Study, With the Number of Items Classified Into Each Outcome

	AHCE	ESDS	ESD2	FTP	HLSU	NLSA	NLS2	NLSC	NLSY	NSHD	TBSS	TLCS	WCFA	YTP	
<i>Physical</i>															
Cardiovascular	10	18	9	2	1	10	2	4	19	23	0	0	0	1	
Development milestones	0	28	2	12	1	9	5	49	1	12	1	10	11	0	
Endocrine	1	5	4	0	1	1	2	1	5	5	0	0	0	0	
Functional autonomy	3	26	20	43	4	8	3	13	20	38	1	8	0	1	
Gastrointestinal	5	16	17	0	0	1	0	1	7	17	0	0	0	0	
General health history	12	50	110	12	28	25	20	47	55	48	2	47	10	1	
Genitourinary	41	21	29	0	0	12	9	4	10	70	0	2	1	1	
Heme-immune	8	4	6	0	2	3	2	2	1	2	1	0	0	0	
Infectious parasitic	0	1	6	1	0	1	0	0	2	6	0	4	0	0	
Metabolic/nutrition	14	28	16	9	5	11	5	14	4	28	4	4	4	0	
Musculoskeletal	8	12	16	3	4	2	0	1	14	9	0	0	0	0	
Nervous	10	30	20	7	7	6	2	5	12	15	0	6	0	1	
Respiratory	6	22	41	4	8	5	2	21	15	31	0	0	0	1	
Sensory	3	56	43	0	3	9	11	8	8	22	2	7	0	0	
Skin	3	18	23	2	1	3	0	0	10	9	0	2	0	0	
Special health needs	0	10	10	0	1	1	5	15	1	19	2	1	0	0	
<i>Well-being</i>															
Accomplishment	1	13	20	4	1	7	17	12	5	5	25	14	1	16	
Engagement	20	15	46	15	7	25	17	83	4	21	22	17	5	63	
Happiness	0	2	1	11	5	3	2	12	3	1	5	2	0	5	
Life satisfaction	0	13	7	19	12	11	5	20	6	20	18	34	1	93	
Meaning/purpose	0	1	3	0	2	4	3	1	0	0	0	0	0	1	
Optimism	1	4	6	5	9	11	20	8	5	1	1	2	0	49	
Perseverance	5	7	8	8	12	4	0	18	7	2	6	11	0	46	
Relations/connectedness	7	24	40	186	55	73	31	107	26	15	26	8	7	53	
<i>Ill-being</i>															
Anxiety	22	51	21	25	3	3	1	9	2	19	5	17	0	16	
Depression	49	19	17	18	6	9	4	9	14	36	3	2	0	6	
Externalizing behavior	37	28	54	109	0	85	49	64	10	1	15	5	20	4	
Negative affect	2	8	15	5	2	2	0	13	6	0	0	6	0	0	
Other psychopathology	15	19	24	3	0	9	2	16	2	8	3	2	3	0	
Stress reactions	1	2	2	2	0	4	0	2	0	5	0	0	0	2	
<i>Health behaviors</i>															
Alcohol use	33	42	32	25	2	27	4	35	53	8	5	4	1	0	
Diet/ nutrition	3	52	36	0	3	44	8	0	4	7	4	1	0	0	
Drug use	53	28	19	30	0	49	5	71	55	0	16	0	2	0	
Other behaviors	3	6	17	0	5	12	3	0	0	7	2	0	2	0	
Physical activity	2	85	7	2	14	15	7	6	5	16	7	7	0	5	
Sexual behavior	23	1	7	6	17	18	19	18	11	7	1	0	4	3	
Sleep	5	16	5	3	4	10	5	3	10	7	3	5	0	0	
Tobacco/ smoking	12	10	12	2	1	14	3	13	13	15	2	1	1	0	
<i>Individual diff.</i>															
Coping style	0	0	0	0	5	0	0	0	0	0	0	1	0	11	
Intelligence	0	8	0	0	0	2	1	21	0	0	2	9	2	10	
Personality	0	17	2	5	1	8	7	36	3	0	14	56	11	9	
Self-esteem	13	2	4	24	3	6	0	32	11	24	10	6	0	28	
Temperament	0	0	0	1	0	1	0	62	0	0	0	8	0	0	
<i>Socio-environmental</i>															
Caregiving	0	7	3	0	0	0	1	1	2	0	3	1	0	0	
Environment	0	40	7	6	0	4	2	69	9	3	1	5	2	32	
Life events	19	24	22	21	8	27	37	31	35	21	11	41	1	2	
Neighborhood	8	1	6	0	1	3	3	5	8	0	4	2	6	0	
Parenting	17	46	30	200	0	40	73	205	17	0	12	13	42	21	
Socio-economic status	14	41	49	24	32	30	98	64	27	23	22	31	23	19	
Victimization/ abuse	14	11	9	2	0	19	11	5	0	0	0	1	6	0	
Total # of coded items	503	988	903	856	276	686	506	1,236	537	626	261	403	166	500	

Note. Numbers indicate how many items from a study were coded into that sub-outcome, at any measurement occasion. See Supplemental Material 1 for outcome definitions and Table 1 for full study names.

There are both strengths and limitations of the TADPOHLS data base. It currently contains 14 studies that can be accessed by researchers, although some are difficult to work with or costly to gain access to. Our data base does not make the data from these studies available; rather, it is designed to classify the information to make it easier for an analyst to know what data are available. Although there are many extant studies that could have been included in the data base, we purposely focused on studies bridging the transition from adolescence into adulthood. The extensive classification scheme used in the data base makes it possible to scale it to many other studies. We hope that additional studies will be added over time, building the resource as a whole to enable collaborative and integrative work.

ILLUSTRATIVE EXAMPLE UTILIZING THE DATA BASE

The data base can be queried to select specific constructs and locate studies with overlapping constructs and items. To illustrate, we examined whether perseverance and optimism measured in adolescence predicted four physical health outcomes measured in adulthood, 15 to 20 years later: self-rated health, physical energy, fatigue, and cardiovascular-related conditions. As an illustration of the potential for the data base as a resource, we used a rudimentary harmonization method to align items. Better methods are still being developed for psychological data, which are much more challenging to collect and harmonize than medical data (Bauer & Hussong, 2009 provides a good example). This limitation should be kept in mind when interpreting the results.

Prior research has suggested associations between positive psychological functioning and better physical health, both cross-sectionally and longitudinally (Diener & Chan, 2011; Howell, Kern, & Lyubomirsky, 2007; Pressman, Gallagher, & Lopez, 2013). Although hundreds of studies have examined this association, most have focused on adults. We aimed to better understand whether this association also pertains to the transition between adolescence and adulthood, using two psychological constructs and four physical health constructs.

Perseverance refers to the tendency to work hard and stick with tasks despite challenges or setbacks. It is a facet of the Big Five personality factor of conscientiousness, which has repeatedly demonstrated small but meaningful associations with health-related outcomes, including better self- and physician-rated health and longer life (e.g., Friedman & Kern, 2014; Kern & Friedman, 2008; Roberts, Lejuez, Krueger, Richards, & Hill, 2014). Optimism refers to the tendency to have

hope and positive expectations for the future, or alternatively as an explanatory style in which good events are seen as internal, stable, and global; and negative events are seen as external, unstable, and specific to the person. Optimism has been related to less reported pain, better physical function, fewer physical symptoms, lower risk of heart disease, and faster recovery from surgery (Boehm & Kubzansky, 2012; Carver, Scheier, & Segerstrom, 2010; Rasmussen, Scheier, & Greenhouse, 2009). We predicted that on average across samples, higher perseverance or optimism would be related to higher levels of energy, better self-rated health, and less cardiovascular disease and fatigue.

Method

Study Selection

We searched the data base for items related to perseverance or optimism during an adolescent assessment (age 13 to 18). We examined items that had been coded into the optimism or perseverance outcomes into the data base (under the positive psychological functioning category), and also examined items in other similar categories, in case of item misspecification. It appeared that items had been correctly specified during the coding process. Items were rated for relevance to our definitions; a few items were excluded as irrelevant (e.g., items assessing self-esteem rather than optimism). If studies had relevant perseverance/optimism items assessed in the adolescent time period, we then examined whether there were physical health items assessed in adulthood. Our final inclusion criteria were: (a) at least one item measuring perseverance and/or optimism in adolescence (age 13–18), and (b) at least one item measuring self-rated health, physical energy, fatigue, or heart conditions in adulthood (age 28–36).

Included Studies

Altogether, four studies had items measuring perseverance and five studies had items measuring optimism. For perseverance, we included the British Birth Cohort Study, 1958 cohort (ESDS), the Family Transition Project (FTP), the National Longitudinal Study of Adolescent Health (NLSA), and the Terman Life Cycle Study (TLCS). For optimism, we included the Family Transition Project (FTP), National Longitudinal Survey of Youth, 1979 Cohort - Children and Youth (NLSC) and 1997 Cohort (NLS2), the National Longitudinal Study of Adolescent Health (NLSA), and the Terman Life Cycle Study (TLCS). Table 1 and Supplemental Material 2 provide sample descriptions, and Supplemental Material 3 provides items included from each study.

Data Analyses

To harmonize the perseverance and optimism items, scores were standardized and averaged to create an overall measure of the construct.² For health outcomes, variables were dichotomized (e.g., self-rated health: poor versus good/very good; heart conditions: present versus absent). In each study, we first computed Spearman rho correlations between perseverance/optimism and each health outcome. We then meta-analytically combined the r effect sizes, using a fixed effects model. To combine effects, we transformed the r s to the Fisher equivalent, $Zr = .5 * \ln((1 + r)/(1 - r))$. Zr s were weighted by sample size (degrees of freedom, calculated as $df = n - 3$ for each study), and then averaged together and 95% confidence intervals were calculated. Values were converted back to r s for presentation purposes (Rosenthal & DiMatteo, 2001).

RESULTS

Table 3 summarizes Spearman rho correlations between perseverance or optimism and physical health, separately in each sample. Across studies, both optimism and perseverance were positively related to self-rated health. Associations varied across the other outcomes. The average correlation and 95% confidence intervals are summarized in the right column. In general, correlations aligned across the individual studies. Both perseverance and optimism were positively associated with self-rated health and were negatively related to fatigue. Optimism was also positively related to physical energy.

DISCUSSION

In this illustrative example, we demonstrated how the data base can be used to identify items representing the same concepts in different studies, and then combined to examine overall effects. Before analyses were performed, items were harmonized, such that analyses in each study were aligned, and the meta-analytic combination was based upon the aligned variables. Adolescent perseverance and optimism were related to better self-reported physical health outcomes 15 to 20 years later. As analyses were limited to the studies within the data base, effects are generalizable only to the

²It is possible that responses differ due to the question wording and/or sample characteristics. IDA allows such differences to be directly tested and included in the model. As the current analysis is an illustrative example, we simply harmonized variables. To fully consider associations between adolescent perseverance and optimism and adult health, more sophisticated approaches to harmonization should be used.

studies included here, but offer support for the importance of optimism and perseverance as protective adolescent characteristics that potentially should be supported and further developed.

It is striking that both optimism and perseverance were predictive of better health outcomes over a 15 to 20 year period, and across diverse samples in terms of geographical location, period in time, and other individual factors. The effect sizes were small in size, and yet were significant when combined across multiple studies and many participants, demonstrating the added power and value of combining data at the individual participant level. Many factors influence variations in physical health outcomes, such that small effects can be practically meaningful (Rosenthal & Rosnow, 2008). The pattern of associations is similar to studies that have linked perseverance, optimism, and related attributes to health outcomes in adults, providing a proof of concept for the utility of the data base and the use of integrative approaches. Subsequent studies can use other variables available in these data sets to examine processes and moderators of these associations.

Two of the samples were very large, while the others were relatively small. The meta-analytic results are mostly determined by the larger samples, such that meta-analysis may not be useful in this situation. Still, smaller studies can complement large nationally representative samples. Large samples often can only include one or two items for a construct, whereas a small study can include richer assessments. For example, the Family Transition Project is an intensive study of rural families, with hundreds of items on parent/child relationships, parenting, and externalizing behaviors (Conger & Conger, 2002; Conger & Elder, 1994). Integrative techniques can be used to link studies, and then be extended to the unique information offered by each study (McArdle, Grimm, Hagemami, Bowles, & Meredith, 2009). Further, when results align across the large and small studies, it provides greater confidence in the overall pattern of findings.

GENERAL DISCUSSION

Developmental psychology has a rich history of studying developmental trajectories, but cross-sectional and longitudinal studies can provide contradictory information. Schaie and others introduced cross-sequential and other study designs for separating age, time of measurement, and cohort effects and understanding different influences on outcomes of interest (e.g., Salthouse & Nesselroade, 2002; Schaie, 1965; Schaie & Strother, 1968). Integrating existing longitudinal studies together may provide a quasi-cross-sequential approach for studying developmental trajectories.

TABLE 3

Prospective Associations Between Adolescent Perseverance or Optimism and Physical Health (Self-Rated Health, Energy, Fatigue, Heart Conditions) for Each Sample (Parallel Analysis) and the Combined Sample (Meta-Analysis of Effect Sizes)

	<i>ESDS</i>		<i>FTP</i>		<i>NLSA</i>		<i>NLS2</i>		<i>NLSC</i>		<i>TLCS</i>		<i>Fixed Effects Meta-analysis</i>				
	N	r	N	r	N	r	N	r	N	r	N	r	K	N	r	r LL	r UL
<i>Perseverance</i>																	
Gender ^a	9,724	.10	334	.09	1,3884	.09					334	.18	4	38,828	.10	.08	.11
Heart conditions ^b	9,013	.00	334	-.03	1,3884	.00					318	-.06	4	38,085	.00	-.01	.01
Energy ^c			334	.03									1	668	.03	-.08	.14
Fatigue ^c	9,017	-.06	334	-.13	1,3883	-.04							3	37,451	-.05	-.06	-.03
Self-rated health ^c	8,977	.09	334	-.02	1,3886	.06					299	.01	4	38,015	.07	.05	.08
<i>Optimism</i>																	
Gender ^a			333	-.04	1,3871	-.05	3,501	.00	838	-.04	334	.00	5	18,877	-.04	-.05	-.02
Heart conditions ^b			333	.01	1,3869	-.01			838	.00	318	-.07	4	15,358	-.01	-.03	.00
Energy ^c			333	-.03			3,043	.07	788	.05			3	4,164	.06	.03	.09
Fatigue ^c			333	-.01	1,3869	-.08							2	14,202	-.07	-.09	-.06
Self-rated health ^c			333	-.08	1,3871	.11	3,500	.02	838	.07	299	.07	5	18,841	.09	.08	.10

^a0 = male, 1 = female; ^b0 = no heart condition, 1 = one or more heart conditions; ^c0 = low, 1 = high.

Note. Gender, perseverance, and engagement were assessed in adolescence; health outcomes were assessed in adulthood, 15–20 years later. Significant correlations ($p < .05$) are bolded. K = number of studies included in meta-analysis calculation; LL = lower 95% confidence interval; UL = upper 95% confidence interval; ESDS = British Birth Cohort Study: 1958 Cohort; FTP = Family Transitions Project; NLSA = National Longitudinal Study of Adolescent Health; NLS2 = National Longitudinal Survey of Youth - 1997 Cohort; NLSC = National Longitudinal Survey of Youth - 1979 Child and Young Adult; TLCS = Terman Life Cycle Study.

We developed the TADPOHLS data base as a resource for researchers who are interested in advancing longitudinal study of human development. Strengths of this resource include providing the opportunity to piece together different cohorts and constructs, and focusing explicitly on studies that followed individuals from adolescence in to adulthood. Our analysis of perseverance, optimism, and physical health provides an illustration of using the TADPOHLS resource.

Studies of psychological well-being and physical health often occur independently. A more holistic and integrated approach involves identifying protective factors that lead to optimal psychological and physical functioning across years or decades, while simultaneously attending to physical, mental, and social components of the individual. Psychological, social, and physical health are interrelated outcomes that are developed and influenced by personality, socioecological context, habits and behaviors, and experiences (Friedman & Kern, 2014). Furthermore, health and psychological research has traditionally focused on negative aspects of risk, atypical development, and disease onset and progression. Although it is certainly important to identify and reduce risk factors, it is also beneficial to identify and strengthen assets that buffer against disease, strengthen an individual's adaptability, and promote thriving. The TADPOHLS data base includes positive psychological characteristics, which can be explored as protective factors from disease.

A growing number of longitudinal studies have examined predictors of adolescent and adult outcomes. For example, in one study, adolescents with high levels

of positive affect and self-esteem reported better overall health, and engaged in fewer risky behaviors across a six-year period (Hoyt, Chase-Lansdale, McDade, & Adam, 2012). Adults with high levels of optimism were at lower risk for developing coronary heart disease across a 10-year period (Kubzansky, Sparrow, Vokonas, & Kawachi, 2001). There are many constructs yet to be investigated, and moderators and processes of such relationships are relatively unknown. By combining multiple studies, developmental trajectories, moderators, and processes impacting such trajectories, and boundary conditions of such associations can potentially be investigated. Both cross-sectional and longitudinal studies make various assumptions that limit conclusions that can be made about development (Schaie, 1965); combining multiple longitudinal studies at the item level provides the potential to test age and cohort-related effects.

At the same time, any attempts to combine data should proceed cautiously. As sample sizes increase, many associations will be statistically significant but not necessarily meaningful. Creating mega-samples simply to reach significance is not useful, but when data are combined directly to test specific theories or to generate new hypotheses, combined data may be useful. In our example, perseverance and optimism were positively related to good health outcomes. This finding aligns with other studies and provides a proof of concept for using studies in the data base. However, our approach was rather rudimentary and conclusions stemming from this analysis are limited. In addition, combining data and increasing sample size potentially limits Type II

errors, highlighting potential associations that should be studied in more detail in subsequent studies. With health outcomes, small effects can be important (Meyer et al., 2001), and large samples are needed to find such effects. By integrating multiple studies, we might uncover important health protective factors that otherwise would be missed with smaller samples.

There are a growing number of collaborations and groups both within and across disciplines, such as the NIH Patient Reported Outcomes Measurement Information Systems (PROMIS; Cella et al., 2007), the NIMH Collaborative Data Synthesis for Adolescent Depression Trials group (CDSADT; Perrino et al., 2013), the eXtending Treatments, Education and Networks in Depression study (xTEND; Allen et al., 2013), Data Aggregation Through Anonymous Summary-statistics from Harmonised Individual level Data bases (DataSHIELD; Jones et al., 2012), and the Grid Enabled Measures Data base (GEM, Moser et al., 2011), among many others. IALSA is an open and growing network, and inspired our work here. The TADPOHLS data base complements these existing resources.

In conclusion, we have developed a resource that contributes toward life span studies of positive psychological and physical health during the transition from adolescence into adulthood. Many of the studies in the data base have already influenced public policy, institutional practices, family, and individual discourse. However, it is clear there is still much to be learned and disseminated to both identify and promote psychological and physical health development in all stages of life.

FUNDING

This research was supported by the Robert Wood Johnson Foundation's Pioneer Portfolio grant, "Exploring the Concepts of Positive Health," and the Klaus J. Jacobs Foundation.

SUPPLEMENTAL MATERIAL

Supplemental material for this article is available on the [publisher's website](#).

REFERENCES

- Allen, J., Inder, K. J., Lewin, T. J., Attia, J. R., Kay-Lambkin, F. J., Baker, A. L., . . . Kelly, B. J. (2013). Integrating and extending cohort studies: Lessons from the extending treatments, education and networks in depression (xTEND) study. *BMC Medical Research Methodology*, *13*, 122–136. doi:10.1186/1471-2288-13-122
- Bath, P. A., Deeg, D., & Poppelaars, J. (2010). The harmonization of longitudinal data: A case study using data from cohort studies in The Netherlands and the United Kingdom. *Ageing and Society*, *30*, 1419–1437. doi:10.1017/S0144686x1000070X
- Bauer, D. J., & Hussong, A. M. (2009). Psychometric approaches for developing commensurate measures across independent studies: Traditional and new models. *Psychological Methods*, *14*, 101–125. doi:10.1037/a0015583
- Block, J. (1995). A contrarian view of the five-factor approach to personality description. *Psychological Bulletin*, *117*, 187–215. doi:10.1037/0033-2909.117.2.187
- Boehm, J. K., & Kubzansky, L. D. (2012). The heart's content: The association between positive psychological well-being and cardiovascular health. *Psychological Bulletin*, *138*, 655–691. doi:10.1037/a0027448
- Carver, C. S., Scheier, M. F., & Segerstrom, S. C. (2010). Optimism. *Clinical Psychology Review*, *30*, 879–889. doi:10.1016/j.cpr.2010.01.006
- Cella, D., Yount, S., Rothrock, N., Gershon, R., Cook, K., Reeve, B., . . . Rose, M. (2007). The patient reported outcomes measurement information system (PROMIS): Progress of an NIH roadmap cooperative group during its first two years. *Medical Care*, *45*, S3–S11. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17443116>
- Conger, R. D., & Conger, K. J. (2002). Resilience in Midwestern families: Selected findings from the first decade of a prospective, longitudinal study. *Journal of Marriage and Family*, *64*, 361–373. doi:10.1111/j.1741-3737.2002.00361.x
- Conger, R. D., & Elder, G. H., Jr. (Eds.). (1994). *Families in troubled times: Adapting to change in rural America. Social institutions and social change*. New York, NY: Aldine de Gruyter.
- Cooper, R., Hardy, R., Sayer, A. A., Ben-Shlomo, Y., Birnie, K., Cooper, C., . . . Kuh, D. (2011). Age and gender differences in physical capability levels from mid-life onwards: The harmonization and meta-analysis of data from eight UK cohort studies. *PLoS ONE*, *6*, e27899. doi:10.1371/journal.pone.0027899
- Curran, P. J., & Hussong, A. M. (2009). Integrative data analysis: The simultaneous analysis of multiple data sets. *Psychological Methods*, *14*, 81–100. doi:10.1037/a0015914
- Diener, E., & Chan, M. (2011). Happy people live longer: Subjective well-being contributes to health and longevity. *Applied Psychology: Health & Well-being*, *3*, 1–43. doi:10.1111/j.1758-0854.2010.01045.x
- Duncan, G. J., Engel, M., Claessens, A., & Dowsett, C. J. (2014). Replication and robustness in developmental research. *Developmental Psychology*, *50*, 2417–2425. doi:10.1037/a0037996
- Friedman, H. S., & Kern, M. L. (2014). Personality, well-being, and health. *Annual Review of Psychology*, *65*, 719–742. doi:10.1146/annurev-psych-010213-115123
- Friedman, H. S., Kern, M. L., Hampson, S. E., & Duckworth, A. L. (2014). A new lifespan approach to conscientiousness and health: Combining the pieces of the causal puzzle. *Developmental Psychology*, *50*, 1377–1389. doi:10.1037/a0030373
- Higgins, J. P. T., & Green, S. (Eds.). (2011). *Cochrane handbook for systematic reviews of interventions* (Version 5.1). *The Cochrane Collaboration*. Retrieved from www.cochrane-handbook.org
- Higgins, J. P. T., Lane, P. W., Anagnostelis, B., Anzures-Cabrera, J., Baker, N. F., Cappelleri, J. C., . . . Whitehead, A. (2013). A tool to assess the quality of a meta-analysis. *Research Synthesis Methods*, *4*, 351–366. doi:10.1002/jrsm.1092
- Hofer, S. M., Berg, S., & Era, P. (2003). Evaluating the interdependence of aging-related changes in visual and auditory acuity, balance, and cognitive functioning. *Psychology and Aging*, *18*, 285–305. doi:10.1037/0882-7974.18.2.285
- Hofer, S. M., & Piccinin, A. M. (2009). Integrative data analysis through coordination of measurement and analysis protocol across independent longitudinal studies. *Psychological Methods*, *14*, 150–164. doi:10.1037/a0015566

- Hofer, S. M., & Piccinin, A. M. (2010). Toward an integrative science of life-span development and aging. *Journal of Gerontology: Psychological Sciences, 65B*, 269–278. doi:10.1093/geronb/gbq017
- Hofer, S. M., & Sliwinski, M. J. (2001). Understanding ageing: An evaluation of research designs for assessing the interdependence of ageing-related changes. *Gerontology, 47*, 341–352. doi:10.1159/000052825
- Hofer, S. M., & Sliwinski, M. J. (2006). Design and analysis of longitudinal studies on aging. In J. E. Birren & K. W. Schaie (Eds.), *Handbook of the psychology of aging* (6th ed., pp. 15–37). Boston, MA: Elsevier Academic Press.
- Howell, R., Kern, M. L., & Lyubomirsky, S. (2007). Health benefits: Meta-analytically determining the impact of well-being on objective health outcomes. *Health Psychology Review, 1*, 83–136. doi:10.1080/17437190701492486
- Hoyt, L. T., Chase-Lansdale, P. L., McDade, T. W., & Adam, E. K. (2012). Positive youth, healthy adults: Does positive wellbeing in adolescence predict better perceived health and fewer risky health behaviors in young adulthood? *Journal of Adolescent Health, 50*, 66–73. doi:10.1016/j.jadohealth.2011.05.002
- Hussong, A. M., Curran, P. J., & Bauer, D. J. (2013). Integrative data analysis in clinical psychology research. *Annual Reviews of Psychology, 9*, 61–89. doi:10.1146/annurev-clinpsy-050212-185522
- Ioannidis, J. P. A. (2012). Why science is not necessarily self-correcting. *Perspective on Psychological Science, 7*, 645–654. doi:10.1177/1745691612464056
- Jones, E. M., Sheehan, N. A., Masca, N., Wallace, S. E., Murtagh, M. J., & Burton, P. R. (2012). DataSHIELD-shared individual-level analysis with sharing the data: A biostatistical perspective. *Norsk Epidemiologi, 21*, 231–239.
- Kern, M. L., & Friedman, H. S. (2008). Do conscientious individuals live longer? A quantitative review. *Health Psychology, 27*, 505–512. doi:10.1037/0278-6133.27.5.505
- Kern, M. L., Hampson, S. E., Goldberg, L. R., & Friedman, H. S. (2014). Integrating prospective longitudinal data: Modeling personality and health in the Terman Life Cycle and Hawaii Longitudinal studies. *Developmental Psychology, 50*, 1390–1406. doi:10.1037/a0030874
- Kubzansky, L. D., Sparrow, D., Vokonas, P., & Kawachi, I. (2001). Is the glass half empty or half full? A prospective study of optimism and coronary heart disease in the Normative Aging Study. *Psychosomatic Medicine, 63*, 910–916. doi:10.1097/00006842-200111000-00009
- Makel, M. C., Plucker, J. A., & Hegarty, B. (2012). Replications in psychology research: How often do they really occur? *Perspectives on Psychological Science, 7*, 537–542. doi:10.1177/1745691612460688
- McArdle, J. J., Grimm, K. J., Hagamami, F., Bowles, R. P., & Meredith, W. (2009). Modeling life-span growth curves of cognition using longitudinal data with multiple samples and changing scale of measurement. *Psychological Methods, 14*, 126–149. doi:10.1037/a0015857
- Meyer, G. J., Finn, S. E., Eyde, L. D., Kay, G. G., Moreland, K. L., Dies, R. R., ... Reed, G. M. (2001). Psychological testing and psychological assessment: A review of evidence and issues. *American Psychologist, 56*, 128–165. doi:10.1037//0003-066x.56.2.128
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G.; The PRISMA Group. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine, 6*(7), e1000097. doi:10.1371/journal.pmed.1000097
- Moser, R. P., Hesse, B. W., Shaikh, A. R., Courtney, P., Morgan, G., Augustson, E., ... Coa, K. (2011). Grid-enabled measures: Using science 2.0 to standardize measures and share data. *American Journal of Preventative Medicine, 40*, S134–S143. doi:10.1016/j.amepre.2011.01.004
- Mroczek, D. K. (2014). Personality plasticity, healthy aging, and interventions. *Developmental Psychology, 50*, 1470–1474. doi:10.1037/a0036028
- Nesselroade, J. R. (2004). Intraindividual variability and short-term change. *Gerontology, 50*, 44–47. doi:10.1159/000074389
- Pashler, H., & Harris, C. R. (2012). Is the replicability crisis overblown? Three arguments examined. *Perspectives on Psychological Science, 7*, 531–536. doi:10.1177/1745691612463401
- Peck, S. C. (2004). *The jingle-jangle jungle of identity: Historical perspective and future directions*. Unpublished manuscript. Retrieved from <http://www.rcgd.isr.umich.edu/garp/articles/peck04.pdf>
- Perrino, T., Howe, G., Sperling, A., Beardslee, W., Sandler, I., Shern, D., ... Brown, C. H. (2013). Advancing science through collaborative data sharing and synthesis. *Perspectives on Psychological Science, 8*, 433–444. doi:10.1177/1745691613491579
- Piccinin, A. M., & Hofer, S. M. (2008). Integrative analysis of longitudinal studies on aging: Collaborative research networks, meta-analysis, and optimizing future studies. In S. M. Hofer & D. F. Alwin (Eds.), *Handbook of cognitive aging: Interdisciplinary perspectives* (pp. 446–476). Thousand Oaks, CA: Sage Publications.
- Pressman, S. D., Gallagher, M. W., & Lopez, S. J. (2013). Is the emotion-health connection a “first-world problem”? *Psychological Science, 24*, 544–549. doi:10.1177/0956797612457382
- Rasmussen, H. N., Scheier, M. F., & Greenhouse, J. B. (2009). Optimism and physical health: A meta-analytic review. *Annals of Behavioral Medicine, 37*, 239–256. doi:10.1007/s12160-009-9111-x
- Roberts, B. W., Lejuez, C., Krueger, R. F., Richards, J. M., & Hill, P. L. (2014). What is conscientiousness and how can it be assessed? *Developmental Psychology, 50*, 1315–1330. doi:10.1037/a0031109
- Rosenthal, R., & DiMatteo, M. R. (2001). Meta-analysis: Recent developments in quantitative methods for literature reviews. *Annual Review of Psychology, 52*, 59–82. doi:10.1146/annurev.psych.52.1.59
- Rosenthal, R., & Rosnow, R. L. (2008). *Essentials of behavioral research: Methods and data analysis* (3rd ed.). Boston, MA: McGraw-Hill.
- Salthouse, T. A., & Nesselroade, J. R. (2002). An examination of the Hofer and Sliwinski evaluation. *Gerontology, 48*, 18–21. doi:10.1159/000048919
- Schaie, K. W. (1965). A general model for the study of developmental problems. *Psychological Bulletin, 64*, 92–107. doi:10.1037/h0022371
- Schaie, K. W., & Strother, C. R. (1968). A cross-sequential study of age changes in cognitive behavior. *Psychological Bulletin, 70*, 671–680. doi:10.1037/h0026811
- Shea, B. J., Grimshaw, J. M., Wells, G. A., Boers, M., Andersson, N., Hamel, C., ... Bouter, L. M. (2007). Development of AMSTAR: A measurement tool to assess the methodological quality of systematic reviews. *BMC Medical Research Methodology, 10*. doi:10.1186/1471-2288-7-10
- Slavich, G. M., & Irwin, M. R. (2014). From stress to inflammation and major depressive disorder: A social signal transduction theory of depression. *Psychological Bulletin, 140*, 774–815. doi:10.1037/a0035302
- Staats, A. W. (1999). Unifying psychology requires new infrastructure, theory, method, and a research agenda. *Review of General Psychology, 3*, 3–13. doi:10.1037/1089-2680.3.1.3
- Stewart, L. A., & Clarke, M. J. (1995). Practical methodology of meta-analyses (overviews) using updated individual patient data. *Statistics in Medicine, 14*, 2057–2079. doi:10.1002/sim.4780141902